



pMSSM SUSY Searches @ 7 TeV





CMS
A Compact Solenoidal Detector for LHC



J.A. Conley, J. S. Gainer, J. L. Hewett, M.-P. Le & TGR
arXiv:1009.2539,1103.1697

T.G. Rizzo

04/13/11



ATLAS & CMS have already made a dent in SUSY space

- **However**, as these searches proceed we need to be **sure** that the analyses **don't miss anything** by **assuming specific SUSY breaking mechanisms** such as mSUGRA, GMSB, AMSB, etc.
- **How do we do this?** There are **several possible approaches**...

Issues:

- The general MSSM is too difficult to study due to the large number of soft SUSY breaking parameters (~ 100).
- Many analyses limited to specific SUSY breaking scenarios having only a few parameters...can we be more general ?

→ Model Generation Assumptions :

- The most general, CP-conserving MSSM with R-parity
 - Minimal Flavor Violation at the TeV scale
 - The lightest neutralino is the LSP & a thermal relic.
 - The first two sfermion generations are degenerate & have negligible Yukawa's.
- These choices mostly control flavor issues producing a fairly general scenario for collider & other studies → the pMSSM

19 pMSSM Parameters

10 sfermion masses: $m_{Q_1}, m_{Q_3}, m_{u_1}, m_{d_1}, m_{u_3}, m_{d_3}, m_{L_1},$
 $m_{L_3}, m_{e_1}, m_{e_3}$

3 gaugino masses: M_1, M_2, M_3

3 tri-linear couplings: A_b, A_t, A_τ

3 Higgs/Higgsino: $\mu, M_A, \tan\beta$

How? Perform 2 Random Scans

Flat Priors

emphasizes moderate masses

$$100 \text{ GeV} \leq m_{\text{sfermions}} \leq 1 \text{ TeV}$$

$$50 \text{ GeV} \leq |M_1, M_2, \mu| \leq 1 \text{ TeV}$$

$$100 \text{ GeV} \leq M_3 \leq 1 \text{ TeV}$$

$$\sim 0.5 M_Z \leq M_A \leq 1 \text{ TeV}$$

$$1 \leq \tan\beta \leq 50$$

$$|A_{t,b,\tau}| \leq 1 \text{ TeV}$$

Log Priors

emphasizes lower masses but **also** extends to higher masses

$$100 \text{ GeV} \leq m_{\text{sfermions}} \leq 3 \text{ TeV}$$

$$10 \text{ GeV} \leq |M_1, M_2, \mu| \leq 3 \text{ TeV}$$

$$100 \text{ GeV} \leq M_3 \leq 3 \text{ TeV}$$

$$\sim 0.5 M_Z \leq M_A \leq 3 \text{ TeV}$$

$$1 \leq \tan\beta \leq 60 \text{ (flat prior)}$$

$$10 \text{ GeV} \leq |A_{t,b,\tau}| \leq 3 \text{ TeV}$$

- **Flat Priors** : 10^7 models scanned , 68422 survive
- **Log Priors** : 2×10^6 models scanned , 2908 survive

→ **Comparison of these two scans will show the prior sensitivity,**

Some Constraints

- **W/Z ratio** **b → s γ**
- **$\Delta(g-2)_\mu$** **$\Gamma(Z \rightarrow \text{invisible})$**
- **Meson-Antimeson Mixing**
- **$B_s \rightarrow \mu\mu$** **$B \rightarrow \tau\nu$**

- **DM density:** $\Omega h^2 < 0.121$. We treat this only as an *upper bound* on the neutralino thermal relic contribution
- **Direct Detection Searches for DM (CDMS, XENON...)**
- **LEP and Tevatron Direct Higgs & SUSY searches :** there are *many* searches & some are quite complicated with many caveats.... These needed to be 'revisited' for the more general case considered here → simulations **limit** model set size (~1 core-century for set generation)

ATLAS SUSY Analyses w/ a Large Model Set

- We passed these points through the ATLAS inclusive MET analyses (@ both 7 & 14 TeV !), designed for mSUGRA , to explore this broader class of models (~150 core-yrs)
- We used the ATLAS SM backgrounds with their associated systematic errors, search analyses/cuts & criterion for SUSY discovery. (→ ATL-PHYS-PUB-2010-010 for 7 TeV)
- We verified that we can approximately reproduce the 7 & 14 TeV ATLAS results for their benchmark mSUGRA models with our analysis techniques for each channel. ..BUT beware of some analysis differences:

ATLAS

ISASUGRA generates spectrum
& sparticle decays

Partial NLO cross sections using
PROSPINO & CTEQ6M

Herwig for fragmentation &
hadronization

GEANT4 for full detector sim

US

SuSpect generates spectra
with SUSY-HIT# for decays



NLO cross section for all 85
processes using PROSPINO**
& CTEQ6.6M

PYTHIA for fragmentation &
hadronization



PGS4-ATLAS for fast detector
simulation

** version w/ negative K-factor errors corrected


version w/o negative QCD corrections, with 1st & 2nd generation fermion masses & other very numerous PS fixes included. e.g., explicit small Δm chargino decays, etc.




7 TeV



M_{eff} distribution for 4-jet, 0 lepton analysis




M_{eff} distribution for 2-jet, 0 lepton analysis




14 TeV

4j

M_{eff} distribution for \hat{M} lepton analysis



M_{eff} distribution for b-jet analysis





→ We do quite well reproducing ATLAS 7 & 14 TeV benchmarks with some small differences due to, e.g., (modified) public code usages & **PGS vs GEANT4**



- The first question: ‘How well do the ATLAS analyses cover the pMSSM model sets?’ More precisely, ‘what fraction of these models can be discovered (or not!) by any of the ATLAS analyses & which ones do best?’
- Then we need to understand **WHY** some models are missed by these analyses even when high luminosities are available

FLAT

Solid=4j, dash=3j, dot=2j final states




Red=20%, green=50%, blue=100% background systematic errors




LOG

Solid=4j, dash=3j, dot=2j final states

Percent Found by (4,3,2)j0l




Percent Found by (4,3,2)jOSDL




Red=20%, green=50%, blue=100% background systematic errors

Percent Found by (4,3,2)j1l



Percent Found by 2jSSDL



What fraction of models are found by **n** analyses @7 TeV assuming, e.g., $\delta B=20\%$?


# anl.	Flat $\mathcal{L}_{0.1}$	Flat \mathcal{L}_1	Flat \mathcal{L}_{10}	Log $\mathcal{L}_{0.1}$	Log \mathcal{L}_1	Log \mathcal{L}_{10}
0	38.172	7.5501	0.9965	63.64	43.988	22.92
1	9.2928	4.1988	0.90862	5.376	4.8674	5.8482
2	8.7432	4.6665	1.6102	3.6687	5.6665	6.0298
3	41.836	59.878	39.573	26.008	34.907	35.38
4	0.65686	4.9257	7.9422	0.25427	2.2158	6.4657
5	0.53472	4.2629	6.7163	0.47221	2.0341	4.8311
6	0.54366	8.5391	13.494	0.32692	3.0875	6.5383
7	0.067026	2.5217	8.9044	0.21794	1.453	4.1773
8	0.062558	1.2288	5.6364	0.036324	0.72648	2.2884
9	0.077452	1.2958	6.548	0	0.58118	2.9422
10	0.013405	0.93241	7.6711	0	0.47221	2.579

→ → SUSY signals usually seen in multiple analyses


How good is the pMSSM coverage @ 7 TeV as the luminosity **evolves** ??

The coverage is quite good for both model sets !

Flat priors



Log priors



- These figures emphasize the importance of decreasing background systematic errors to obtain good **pMSSM** model coverage. For FLAT priors we see that, e.g.,

$L=5(10) \text{ fb}^{-1}$ and $\delta B=100\%$ is 'equivalent' to

$L=0.65(1.4) \text{ fb}^{-1}$ and $\delta B=50\%$ ($x \sim 7$) OR to

$L=0.20(0.39) \text{ fb}^{-1}$ and $\delta B=20\%$ ($x \sim 25$) !!

This effect is **less dramatic** for the LOG case due to the potentially heavier & possibly compressed mass spectrum

ATLAS pMSSM Model Coverage*

RIGHT NOW for $\sim 35 \text{ pb}^{-1}$ @ 7 TeV

<u>δB</u> :	<u>100%</u>	<u>50%</u>	<u>20%</u>
FLAT:	16%	29%	39%
LOG:	11%	20%	27%

Wow! This is actually quite impressive as these LHC SUSY searches are **just beginning** !

* Fraction of models that **SHOULD** have been found but weren't if all ATLAS analyses were performed as stated

Search 'effectiveness': If a model is found by **only 1** analysis which one is it??

Analysis	Flat $\mathcal{L}_{0.1}$	Flat \mathcal{L}_1	Flat \mathcal{L}_{10}	Log $\mathcal{L}_{0.1}$	Log \mathcal{L}_1	Log \mathcal{L}_{10}
4j0l	71.037	63.533	59.18	75.676	63.433	41.615
3j0l	1.154	11.493	18.689	1.3514	11.94	21.118
2j0l	26.206	13.799	4.4262	20.27	15.672	12.422
4j1l	0.30454	4.6116	6.5574	0	5.9701	7.4534
3j1l	0.096169	0.81589	0.98361	0	0	0.62112
2j1l	0.080141	1.8801	4.0984	0	0	6.2112
4jOSDL	0.048085	0	0	0	0.74627	0
3jOSDL	0.032056	1.6318	0.32787	0	0	0.62112
2jOSDL	0.99375	1.6673	0.4918	1.3514	1.4925	1.8634
2jSSDL	0.048085	0.56758	5.2459	1.3514	0.74627	8.0745

$\delta B = 20\%$

→ → again, 4j0l is the most powerful analysis...


The Undiscovered SUSY

Why Do Models Get Missed by ATLAS?

The most obvious things to look at **first** are :


- **small signal rates due to suppressed σ 's**
- **which can be correlated with large sparticle masses**
- **small mass splittings w/ the LSP (compressed spectra)**
- **decay chains ending in stable charged sparticles**

σ_{gg} σ_{SS} vs gluino mass and lightest squark mass




σ 's: Squark & gluino production cross sections @ 7 TeV cover a very **wide range** & are correlated with the **search significance**. But there are models with $\sigma \sim 30$ pb that are **missed by all ATLAS analyses** while others with σ below ~ 100 fb **are found**.

Significance vs σ_{sQCD}




σ_{gg} production vs σ_{SS} production




Soft jets & leptons

Both 7 & 14 TeV models can be missed due to **small mass splittings** between squarks and/or gluinos and the LSP \rightarrow softer jets or leptons not passing cuts. **ISR helps in some cases...**

Models that fail all analyses for flat priors, 1 fb^{-1}




4j0l analysis for flat priors, 1 fb^{-1}



7 TeV

4j0l analysis for flat priors, 1 fb^{-1}




For small mass splittings w/ the LSP a smaller fraction of events will pass analysis cuts

of evts passing cuts
total generated

Red=squark pairs
Green=gluino pairs



But as seen on the previous slide tiny efficiencies can be compensated for by huge σ 's !



4j0l

Mass Splitting with the LSP


Missed vs Found Model Comparisons




- 38036 (~2.5 pb) fails while 47772 (~1.7 pb) passes all nj0l
- u_R lighter (~500 vs ~635 GeV) & produces larger σ in 38036 but decays ~75% to $j+\text{MET}$ in both models
- BUT due to the Δm w/ LSP difference (\rightarrow eff ~13% vs ~3.5%) 38036 fails to have a large enough rate after cuts
Efficiencies win over cross sections !

Missed vs Found Model Comparisons


34847-21089-JET1_{trigger}




34847-21089-LEP1_{trigger}



M (GeV)



M (GeV)




What went wrong ??



- 21089 ($\sigma \sim 4.6\text{pb}$) & 34847 ($\sigma \sim 3.3\text{pb}$) yet both models fail n_j0l due to smallish Δm 's. BUT 34847 is seen in the lower background channels (3,4) $j1l$
- In 34847, u_R cascades to the LSP via χ_2^0 & the chargino producing leptons via W emission. The LSP is mostly a wino in this case.
- In 21089, however, u_R can only decay to the lighter \sim Higgsino triplet which is sufficiently degenerate as to be incapable of producing high p_T leptons
- Note that the jets in both u_R decays have similar p_T 's

Missed vs Found Model Comparisons

8944-21089-JET1-Trigger



8944-21089-LEP1-Trigger




What went wrong ??




- **8944** seen in (3,4)OSDL while **21089** is completely missed
nj0l fail due to spectrum compression but with very similar colored sparticle total $\sigma = (3.4, 4.6)$ pb
- models have similar gaugino sectors w/ $\chi_{1,2}^0$ Higgsino-like & χ_3^0 bino-like
- χ_3^0 can decay thru sleptons to produce OSDL + MET
- However in **8944**, the gluino is heavier than d_R so that d_R can decay to χ_3^0
- **But in 21089**, the gluino is lighter than u_R so that it decays into **the gluino** & not **the bino** so NO leptons

Missed vs Found Model Comparisons

9781-20875-LEP1-Trigger



9781-20875-LEP2-Trigger




What went wrong ??




- **9781** seen in 2jSSDL while **20875** is completely missed
nj0l fail due to spectrum compression but with very similar colored sparticle total $\sigma = (1.1, 1.3)$ pb
- Both models have **highly mixed** neutralinos & charginos w/ a relatively compressed spectrum
- **In model 9781**, u_R can decay to $j + \text{leptons} + \text{MET}$ via the **bino** part of χ_2^0 through intermediate e, μ sleptons
- **But in 20875**, these sleptons are **too heavy** to allow for decay on-shell & only **staus** are accessible. The resulting leptons from the taus **are too soft** to pass analysis cuts

Missed vs Found Model Comparisons

68329-10959-JET1_Ttrigger





68329-10959-MET_Ttrigger



What went wrong ??



- **68329** passes 4j0l ($\sigma \sim 4.6$ pb) while **10959** ($\sigma \sim 6.0$ pb) fails all
- In 68329, d_R decays to $j+\text{MET}$ ($B \sim 95\%$) since the gluino is **only ~ 3 GeV lighter**. The gluino decays to the LSP via the sbottom ($B \sim 100\%$) with **a $\Delta m \sim 150$ GeV mass splitting**. The LSP is bino-like in this model
- In 10959, d_R decays via the **~ 107 GeV lighter gluino ($B \sim 99\%$)** and the gluino decays (**with $\Delta m \sim 40$ GeV**) through sbottom & 2nd neutralino to the (wino-like) LSP (**with $\Delta m \sim 60$ GeV**).
- Raising the LSP & b_1 masses in 68239 **by 50 GeV** (the 2nd set of curves) induces **failure** due to the new gluino decay path

Missed vs Found Model Comparisons



- 13900 & 65778 have **heavy spectra & well-mixed gauginos** w/ $\sigma \sim 0.36(0.22)$ pb, too small for nj0l but 65778 seen in 4j1l
- In 13900 the gluino decays to sbottoms & stops while u_R goes mostly to the LSP, so no leptons
- In 65778, $(d,u)_R$ decay to $j+\chi_{2,4}^0$, then to $W\chi_1^\pm$ w/ $B\sim 75\%$ & $\Delta m\sim 160-270$ GeV, producing a subsequent lepton

A 14 TeV Example:



Failed model 43704(process-partonicXS-fullXS-frac.diff)				Sister model 63170			
62	591.6537	552.6714	0.0705342	62	554.1683	598.2279	-0.0736501
63	919.5316	1007.283	-0.0871171	63	1136.412	1115.883	0.0183972
68	1689.407	2207.448	-0.234679	68	1574.955	2111.774	-0.254203
69	4117.824	4558.5	-0.0966714	69	4469.741	4868.156	-0.0818411

#Cut	lepton-pt	num-leps	MET	hardest jet	Meff-4	Meff-3	Meff-2	Sum-4jet-pt	Sum-3jet-pt	Sum-2jet-pt
43704	46.50313	0.3305726	114.8049	424.9652	1070.408	996.6819	859.0967	893.2752	819.5494	681.9642
63170	74.5432	0.3209754	200.8012	368.0755	1090.669	1005.495	867.3606	819.9918	734.8182	596.6838

What went wrong ??

In 43704: gluinos $\rightarrow d_R \rightarrow \chi_2^0 \rightarrow W +$ 'stable' chargino ($\sim 100\%$)
(Zanesville, OH) as the χ_2^0 -LSP mass splitting is ~ 91 GeV


In 63170: gluinos $\rightarrow u_R \rightarrow \chi_2^0 \rightarrow Z/h +$ LSP ($\sim 30\%$) as the
(St. Louis, MO) χ_2^0 -LSP mass splitting is larger ~ 198 GeV

- Again: a small spectrum change can have a large effect on the signal observability!
- \rightarrow Searches for stable charged particles in complex cascades may fill in some gaps as they are common in our model sets


'Stable' Charged Particles in Cascades

→ Mostly long-lived charginos produced in long decay chains

~84% of these χ_1^\pm with $c\tau > 20\text{m}$ have $\sigma_B > 10\text{ fb}$ @ 7 TeV




Unboosted Minimum Decay Length



Estimated σ_B

Impact of Higgs Searches

Searches for the various components of the SUSY Higgs sector also can lead to very important constraints on SUSY parameter space.



So far with $\sim 35 \text{ pb}^{-1}$ these searches have excluded only 4 of our models (due to the existing strong flavor constraints) but these searches are just beginning ..


Summary & Conclusions

- ATLAS searches at **both 7 & 14 TeV** (& **any value** in between) with $\sim 10 \text{ fb}^{-1}$ will do **quite well** at discovering or excluding most of the **FLAT** pMSSM models & **not at all badly** with the **LOG** prior set
- **With $\sim 35 \text{ pb}^{-1}$, a reasonable fraction of this model space has already been 'covered' !**
- **Reducing SM background uncertainties is quite important in enhancing model coverage..**
- **Models 'missed' due to either compressed spectra *or* because of low MET cascades ending in 'stable' charginos *or*...**
There are actually MANY reasons that models are missed.


Summary & Conclusions (cont.)

- Searches in other channel, e.g., **stable charged particles & Higgs**, will play an important role in covering the pMSSM parameter space
- Quite commonly small changes **in the sparticle spectrum can lead to very significant changes in signal rates & will then substantially alter the chances for SUSY discovery**

BACKUP SLIDES




Models that fail all analyses for flat priors, 10 fb^{-1}




Fine-Tuning SUSY ?

- It is often claimed that if the LHC (@7 TeV) does not find anything then SUSY must be VERY fine-tuned & so 'less likely'. Is this true for the pMSSM??

Fine-tuning for models that fail all analyses for FLAT priors and 50% error





Fine-tuning for models that fail all analyses for FLAT priors and 50% error




→ Models w/ low tuning do appear to ‘suffer’ more than those w/ larger values from null SUSY searches

- The amount of fine tuning in the LOG prior set is somewhat less influenced by null ATLAS searches due to spectrum differences , i.e., compression plus mass stretch-out






- How many signal events do we need to reach $S=5$?
Depends on the M_{eff} 'cut' which is now 'optimized' @ 7 TeV

N_S required to get 5σ discovery with various M_{eff} cuts for $nj0l$







• The size of the background systematic error can play a very significant role in the pMSSM model coverage especially for $n_j(0,1)$...




Survivor Spectra : FLAT


g Mass Distribution for FLAT models failed for 50% error

 d_L Mass Distribution for FLAT models failed for 50% error d_R Mass Distribution for FLAT models failed for 50% error u_L Mass Distribution for FLAT models failed for 50% error


u_R Mass Distribution for FLAT models failed for 50% error




t_1 Mass Distribution for FLAT models failed for 50% error







b_1 Mass Distribution for FLAT models failed for 50% error



τ_1 Mass Distribution for FLAT models failed for 50% error



χ_1^0 Mass Distribution for FLAT models failed for 50% error χ_1^+ Mass Distribution for FLAT models failed for 50% error e_L Mass Distribution for FLAT models failed for 50% error e_R Mass Distribution for FLAT models failed for 50% error

Aside: How many models remain missing in the 'best' case as the minimum requirements of 'S=5' for all searches is weakened?

