

The Destruction of Black Holes

arXiv:1207.3342

How rapidly can a Schwarzschild black hole be destroyed?

How rapidly can a Schwarzschild black hole be destroyed?

| Method | Lifetime |
|---------|----------|
| Hawking | M^3 |

How rapidly can a Schwarzschild black hole be destroyed?

| Method | Lifetime |
|---------|-------------------|
| Hawking | $G^2 M^3 / \hbar$ |

How rapidly can a Schwarzschild black hole be destroyed?

| Method | Lifetime |
|-------------|-------------------|
| Hawking | $G^2 M^3 / \hbar$ |
| $\hbar = 0$ | ∞ |

How rapidly can a Schwarzschild black hole be destroyed?

| Method | Lifetime |
|-------------|-------------------|
| Hawking | $G^2 M^3 / \hbar$ |
| $\hbar = 0$ | ∞ |
| Mining?? | $\alpha^{-2} M$ |

How rapidly can a Schwarzschild black hole be destroyed?

| Method | Lifetime |
|-------------|-------------------|
| Hawking | $G^2 M^3 / \hbar$ |
| $\hbar = 0$ | ∞ |
| Mining?? | $\alpha^{-2} M$ |
| Mining | μM^3 |


How rapidly can a Schwarzschild black hole be destroyed?

| Method | Lifetime | Restrictions |
|-------------|-------------------|--------------|
| Hawking | $G^2 M^3 / \hbar$ | • WEC/NEC |
| $\hbar = 0$ | ∞ | |
| Mining?? | $\alpha^{-2} M$ | |
| Mining | μM^3 | |

How rapidly can a Schwarzschild black hole be destroyed?


| Method | Lifetime | Restrictions |
|-------------|-------------------|----------------|
| Hawking | $G^2 M^3 / \hbar$ | • WEC/NEC |
| $\hbar = 0$ | ∞ | $M \gg M_{Pl}$ |
| Mining?? | $\alpha^{-2} M$ | |
| Mining | μM^3 | |

How rapidly can a Schwarzschild black hole be destroyed?

| Method | Lifetime | Restrictions |
|-------------|-------------------|---|
| Hawking | $G^2 M^3 / \hbar$ | <ul style="list-style-type: none">• WEC/NEC |
| $\hbar = 0$ | ∞ | $M \gg M_{Pl}$ |
| Mining?? | $\alpha^{-2} M$ |  |
| Mining | μM^3 | |

$$ds^2 = -\chi^2 dt^2 + \frac{dr^2}{\chi^2} + r^2 d\Omega^2$$


$$\chi \equiv \sqrt{1 - \frac{2M}{r}}$$

$$ds^2 = -\chi^2 dt^2 + \frac{dr^2}{\chi^2} + r^2 d\Omega^2$$


$\chi \ll 1$

$$\chi \equiv \sqrt{1 - \frac{2M}{r}}$$

$$ds^2 = -\chi^2 dt^2 + 4M^2 d\chi^2 + dx^2 + dy^2$$

$$\chi = 0$$


BLACK HOLE

$$ds^2 = -\chi^2 dt^2 + \frac{dr^2}{\chi^2} + r^2 d\Omega^2$$

$$\chi \equiv \sqrt{1 - \frac{2M}{r}}$$

$\chi \ll 1$

$$ds^2 = -\chi^2 dt^2 + 4M^2 d\chi^2 + dx^2 + dy^2$$

χM is:

- distance to horizon
- redshift
- doubling

$\chi = 0$

BLACK HOLE

$$ds^2 = -\chi^2 dt^2 + \frac{dr^2}{\chi^2} + r^2 d\Omega^2$$

$$\chi \equiv \sqrt{1 - \frac{2M}{r}}$$


$$ds^2 = -\chi^2 dt^2 + 4M^2 d\chi^2 + dx^2 + dy^2$$

χM is:

- distance to horizon
- redshift
- doubling

$$(\text{Temp})^{-1}$$

$$\lambda$$



$$ds^2 = -\chi^2 dt^2 + \frac{dr^2}{\chi^2} + r^2 d\Omega^2$$

$$\chi \equiv \sqrt{1 - \frac{2M}{r}}$$

$\chi \ll 1$


$$ds^2 = -\chi^2 dt^2 + 4M^2 d\chi^2 + dx^2 + dy^2$$

χM is:

- distance to horizon
- redshift
- doubling

$(\text{Temp})^{-1}$

λ



$$ds^2 = -\chi^2 dt^2 + \frac{dr^2}{\chi^2} + r^2 d\Omega^2$$

$$\chi \equiv \sqrt{1 - \frac{2M}{r}}$$

$$ds^2 = -\chi^2 dt^2 + 4M^2 d\chi^2 + dx^2 + dy^2$$

χM is:

- distance to horizon
- redshift doubling

$$(\text{Temp})^{-1}$$

$$\lambda$$

